



TITLE:

# On examples of $\mathbb{R}$ -holonomic complexes

AUTHOR(S):

KAWAI, Takahiro

---

CITATION:

KAWAI, Takahiro. On examples of  $\mathbb{R}$ -holonomic complexes.  
数理解析研究所講究録 1993, 854: 20-21

ISSUE DATE:

1993-11

URL:

<http://hdl.handle.net/2433/83751>

RIGHT:

## On examples of $\mathbb{R}$ -holonomic complexes

Takahiro KAWAI (河合 隆久)

Research Institute for Mathematical Sciences

Kyoto University

Kyoto, 606-01 JAPAN

Finding out and analyzing “meaningful” examples of  $\mathbb{R}$ -holonomic complexes (cf. [SKK2]) seems to be one of the most challenging problems in “microlocal analysis in the future”. Examples so far known are related to the  $\theta$ -zerovalues, and they are constructed with the help of the so-called “Jacobi structure”, that is, a set  $\{p_j\}_{1 \leq j, k \leq 2n}$  of microdifferential operators of order  $< 1$  that satisfy

$$[p_j, p_k] = 2\pi\sqrt{-1} e_{jk} \quad (1 \leq j, k \leq 2n)$$

with  $(e_{jk})_{1 \leq j, k \leq 2n}$  being a non-degenerate matrix whose entries are all integers; the infinite order system we are interested in is then constructed as  $(\exp p_j - 1)u = 0$  ( $j = 1, \dots, 2n$ ), roughly speaking. (Cf. [S]). In order to construct a Jacobi structure making use of matrices of differential operators, we usually need to consider some auxiliary systems. (Cf. [SKK1]. Actually the condition on the order of  $p_j$  also becomes somewhat more delicate;  $\text{ord}(\sum c_j p_j) < 1$  for any  $c = (c_1, \dots, c_{2n}) \in \mathbb{C}^{2n}$  is the one employed in [SKK1].) An example of this sort is explicitly written down in [KKT] (for  $n = 2$ ), and detailed analysis of the example is given there; as is known by a general result on  $\mathbb{R}$ -holonomic complexes, the  $\mathcal{O}$ -solution complex of the system in question is  $\mathbb{R}$ -constructible. Still more important is the fact that we can determine its structure explicitly ([KKT], §3); in particular, its first cohomology group is a locally constant sheaf of rank 1 on some stratum, and it has a non-trivial “monodromic” structure. As the system discussed there can be regarded as

the counterpart of the de Rham system in the category of  $\mathbb{R}$ -holonomic complexes (cf. [K2] §3.5), I dare say the  $\mathbb{R}$ -holonomic complexes studied in [K1] (for  $n = 1$ ) and in [KKT] (for  $n = 2$ ) should be the starting point of a concrete study of  $\mathbb{R}$ -holonomic complexes.

### References

- [KKT] Kashiwara, M., T. Kawai and Y. Takei: The structure of cohomology groups associated with the theta-zerovalues, Proc. of the Conference "Geometrical and Algebraical Aspects in Several Complex Variables", Editel (Iatary), pp.169-189, 1991.
- [K1] Kawai, T.: An example of a complex of linear differential operators of infinite order, Proc. Japan Acad. Ser. A, **59** (1983), 113-115.
- [K2] ———: Systems of linear differential equations of infinite order —an aspect of infinite analysis—, Theta Functions, Bowdoin 1987, Proc. Symp. in Pure Math., **49** (1989), Part 1, Amer. Math. Soc., pp.3-17.
- [S] Sato, M.: Pseudo-differential equations and theta functions, Astérisque, **2 et 3** (1973), 286-291.
- [SKK1] Sato, M., M. Kashiwara and T. Kawai: Linear differential equations of infinite order and theta functions, Adv. in Math., **47** (1983), 300-325.
- [SKK2] ———: Microlocal analysis of theta functions, Advanced Studies in Pure Math., **4** (1984), 267-289.